

Table 5.15a
Unadjusted Patient Survival by Year of Transplant at 3 Months, 1 Year,
3 Years, 5 Years and 10 Years
Deceased Donor non-ECD Kidney Transplants

		3 Months		1 Year		3 Years		5 Years		10 Years	
		Surv.	Std. Err.	Surv.	Std. Err.	Surv.	Std. Err.	Surv.	Std. Err.	Surv.	Std. Err.
Trans- #	Trans- plants										
1987	1,355	95.9%	0.5%	92.3%	0.7%	84.7%	1.0%	76.8%	1.1%	58.2%	1.3%
1988	5,782	95.9%	0.3%	92.2%	0.4%	85.4%	0.5%	78.0%	0.5%	58.6%	0.6%
1989	5,474	96.0%	0.3%	93.0%	0.3%	86.6%	0.5%	79.9%	0.5%	60.3%	0.7%
1990	5,841	96.4%	0.2%	93.3%	0.3%	87.6%	0.4%	80.8%	0.5%	60.4%	0.6%
1991	5,754	97.4%	0.2%	94.6%	0.3%	88.8%	0.4%	81.1%	0.5%	60.1%	0.6%
1992	5,640	97.0%	0.2%	94.1%	0.3%	88.1%	0.4%	80.0%	0.5%	58.6%	0.7%
1993	6,020	97.4%	0.2%	94.5%	0.3%	88.6%	0.4%	81.3%	0.5%	61.1%	0.6%
1994	5,746	97.7%	0.2%	94.5%	0.3%	88.7%	0.4%	81.4%	0.5%	60.9%	0.6%
1995	5,646	97.3%	0.2%	94.7%	0.3%	89.5%	0.4%	82.2%	0.5%	63.0%	0.6%
1996	5,583	97.5%	0.2%	95.2%	0.3%	89.3%	0.4%	82.4%	0.5%	63.1%	0.6%
1997	5,553	97.4%	0.2%	94.9%	0.3%	89.9%	0.4%	83.0%	0.5%	63.2%	0.6%
1998	5,719	98.0%	0.2%	95.7%	0.3%	90.3%	0.4%	83.9%	0.5%	+	+
1999	5,754	97.4%	0.2%	94.8%	0.3%	89.6%	0.4%	83.0%	0.5%	+	+
2000	5,734	97.7%	0.2%	94.5%	0.3%	89.2%	0.4%	82.2%	0.5%	+	+
2001	5,868	97.9%	0.2%	95.5%	0.3%	89.1%	0.4%	82.1%	0.5%	+	+
2002	5,989	97.8%	0.2%	95.4%	0.3%	89.5%	0.4%	83.1%	0.5%	+	+
2003	5,976	98.1%	0.2%	95.5%	0.3%	90.0%	0.4%	+	+	+	+
2004	6,386	98.3%	0.2%	95.9%	0.2%	90.6%	0.4%	+	+	+	+
2005	6,692	98.3%	0.2%	95.6%	0.3%	+	+	+	+	+	+
2006	7,240	98.3%	0.2%	96.1%	0.2%	+	+	+	+	+	+

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(+) = Values not determined due to insufficient follow-up.

Multi-organ transplants are excluded for kidney, pancreas, heart, lung, and liver survival rates.

Intestine survival includes intestine only and intestine with liver and/or pancreas transplants.

Living donor transplants excluded unless explicitly listed. Heterotopic heart and liver transplants excluded.

Patient survival follows patients from first transplant of this type until death. Graft survival follows individual transplants until graft failure. Counts for patient and graft survival are different because a patient may have more than one transplant for a type organ.

See Technical Notes for complete discussion of patient and graft survival methods. See Technical Notes for details of patient and graft survival methods.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.